

Toric Fano varieties associated to graph cubeahedra

Yusuke Suyama (Osaka University)

Algebraic and Geometric Combinatorics on Lattice Polytopes

August 2, 2018

Table of Contents

- 1 Toric varieties and fans
- 2 Graph cubeahedra
- 3 Toric Fano varieties
- 4 Main results

1. Toric varieties and fans

Definition

An n -dimensional **toric variety** is a normal algebraic variety X over \mathbb{C} containing $(\mathbb{C}^*)^n$ as an open dense subset, s.t. the natural action $(\mathbb{C}^*)^n \curvearrowright (\mathbb{C}^*)^n$ extends to an action on X .

Examples

$(\mathbb{C}^*)^n, \mathbb{C}^n, \mathbb{P}^n$ are toric varieties.

Definition

Two toric varieties X and X' are said to be **isomorphic** if there exists an isomorphism $f : X \rightarrow X'$ satisfying the following conditions:

- f induces an isomorphism of algebraic tori $f' : (\mathbb{C}^*)^n \rightarrow (\mathbb{C}^*)^n$.
- f is equivariant with respect to f' , i.e., $f(tx) = f'(t)f(x)$ for any $t \in (\mathbb{C}^*)^n$ and $x \in X$.

Definition

A **rational strongly convex polyhedral cone** is a cone $\sigma \subset \mathbb{R}^n$ generated by finitely many vectors in \mathbb{Z}^n which does not contain any non-zero linear subspace of \mathbb{R}^n . A **fan** in \mathbb{R}^n is a non-empty finite set Δ of such cones satisfying the following conditions:

- If $\sigma \in \Delta$, then each face of σ is in Δ .
- If $\sigma, \tau \in \Delta$, then $\sigma \cap \tau$ is a face of each.

Definition

Two fans Δ and Δ' in \mathbb{R}^n are said to be **isomorphic** if there exists an automorphism of \mathbb{Z}^n that induces a bijection $\Delta \rightarrow \Delta'$.

Theorem

$$\{\text{fans in } \mathbb{R}^n\}/(\text{isom.}) \xleftrightarrow{1:1} \{n\text{-dimensional toric varieties}\}/(\text{isom.}), \\ \Delta \mapsto X(\Delta).$$

We construct a toric variety $X(\Delta)$ from a fan Δ .

Step 1 (affine toric varieties)

For each $\sigma \in \Delta$, we construct an affine toric variety U_σ .

- $\sigma^\vee = \{u \in \mathbb{R}^n \mid \langle u, v \rangle \geq 0 \ \forall v \in \sigma\}$: the **dual** of σ .
- $\sigma^\vee \cap \mathbb{Z}^n$ is a commutative monoid.
- The monoid ring $\mathbb{C}[\sigma^\vee \cap \mathbb{Z}^n]$ is a finitely generated integral domain over \mathbb{C} . So we put $U_\sigma = \text{Spec} \mathbb{C}[\sigma^\vee \cap \mathbb{Z}^n]$.

Step 2 (gluing)

Let τ be a face of σ and let $\tau \rightarrow \sigma$ be the inclusion.



- \leadsto a monoid homomorphism $\sigma^\vee \cap \mathbb{Z}^n \rightarrow \tau^\vee \cap \mathbb{Z}^n$.
- \leadsto an open immersion $U_\tau \rightarrow U_\sigma$.
- Gluing $\{U_\sigma \mid \sigma \in \Delta\}$, we obtain the toric variety $X(\Delta)$.

Example

$$\sigma = \mathbb{R}_{\geq 0}(2\mathbf{e}_1 - \mathbf{e}_2) + \mathbb{R}_{\geq 0}\mathbf{e}_2 \subset \mathbb{R}^2$$


$$\leadsto \sigma^\vee = \mathbb{R}_{\geq 0}\mathbf{e}_1 + \mathbb{R}_{\geq 0}(\mathbf{e}_1 + 2\mathbf{e}_2)$$

$$\leadsto \sigma^\vee \cap \mathbb{Z}^2 = \mathbb{Z}_{\geq 0}\mathbf{e}_1 + \mathbb{Z}_{\geq 0}(\mathbf{e}_1 + \mathbf{e}_2) + \mathbb{Z}_{\geq 0}(\mathbf{e}_1 + 2\mathbf{e}_2).$$




$\mathbb{C}[\sigma^\vee \cap \mathbb{Z}^2] = \mathbb{C}[X, XY, XY^2] = \mathbb{C}[U, V, W]/(UW - V^2)$. Therefore $U_\sigma = \text{Spec} \mathbb{C}[U, V, W]/(UW - V^2)$.


Examples ($n = 2$)



\mathbb{P}^2



$\mathbb{P}^1 \times \mathbb{P}^1$



Hirzebruch surface F_1

Δ : a fan in \mathbb{R}^n .

Definition

- Δ is **nonsingular** \Leftrightarrow every cone of Δ is generated by a part of a basis for \mathbb{Z}^n .
- Δ is **complete** $\Leftrightarrow \bigcup_{\sigma \in \Delta} \sigma = \mathbb{R}^n$.

Fact

- $X(\Delta)$ is nonsingular $\Leftrightarrow \Delta$ is nonsingular.
- $X(\Delta)$ is complete $\Leftrightarrow \Delta$ is complete.

2. Graph cubeahedra

Let G be a **finite simple graph**, that is, a finite graph with no loops and no multiple edges.

- $V(G) = \{1, \dots, n\}$: the node set.
- $E(G)$: the edge set.


Definition

For $I \subset V(G)$, we define the **induced subgraph** $G|_I$ by

$$V(G|_I) = I, \quad E(G|_I) = \{\{v, w\} \in E(G) \mid v, w \in I\}.$$

Let \square^n be an n -cube whose facets are labeled by $1, \dots, n$ and $\bar{1}, \dots, \bar{n}$, where the two facets labeled by i and \bar{i} are on opposite sides.


Then every face of \square^n is labeled by a subset $I \subset \{1, \dots, n, \bar{1}, \dots, \bar{n}\}$ such that $I \cap \{1, \dots, n\}$ and $\{i \in \{1, \dots, n\} \mid \bar{i} \in I\}$ are disjoint. The face corresponding to I is the intersection of the facets labeled by the elements of I .



Let $\mathcal{I}_G = \{I \subset V(G) \mid I \neq \emptyset, G|_I \text{ is connected}\}$. We call \mathcal{I}_G the **graphical building set** of G .

Example

Let P_3 be a path with three nodes.



Then $\mathcal{I}_{P_3} = \{\{1\}, \{2\}, \{3\}, \{1, 2\}, \{2, 3\}, \{1, 2, 3\}\}$.

Definition (Devadoss–Heath–Vipismakul, 2011)

The **graph cubeahedron** \square_G is obtained from \square^n by truncating the faces labeled by the elements of \mathcal{I}_G in increasing order of dimension.



Example

Let P_n be a path with n nodes. Then

$$\mathcal{I}_{P_2} = \{\{1\}, \{2\}, \{1, 2\}\},$$

$$\mathcal{I}_{P_3} = \{\{1\}, \{2\}, \{3\}, \{1, 2\}, \{2, 3\}, \{1, 2, 3\}\}$$

and we have the graph cubeahedra \square_{P_2} and \square_{P_3} in the following figures:



We have a one-to-one correspondence

$$\begin{aligned} \mathcal{I}_G \cup \{\{\bar{1}\}, \dots, \{\bar{n}\}\} &\xleftrightarrow{1:1} \{\text{facets of } \square_G\}, \\ I &\mapsto F_I. \end{aligned}$$

Theorem (Devadoss–Heath–Vipismakul, 2011)

Let G be a finite simple graph. Then the two facets F_I and F_J of the graph cubeahedron \square_G intersect iff one of the following holds:


- $I, J \in \mathcal{I}_G$ and we have either $I \subset J$ or $J \subset I$ or $I \cup J \notin \mathcal{I}_G$.
- One of I and J , say I , is in \mathcal{I}_G and $J = \{\bar{j}\} \quad \exists j \in \{1, \dots, n\} \setminus I$.
- $I = \{\bar{i}\}$ and $J = \{\bar{j}\}$ for some $i, j \in \{1, \dots, n\}$.

Furthermore, \square_G is a flag polytope.

We can realize \square_G as a smooth polytope (or a Delzant polytope) such that for any $I \in \mathcal{I}_G \cup \{\{\bar{1}\}, \dots, \{\bar{n}\}\}$, the outward-pointing primitive normal vector e_I of F_I is

$$e_I = \begin{cases} \sum_{i \in I} e_i & (I \in \mathcal{I}_G), \\ -e_i & (I = \{\bar{i}\}, i \in \{1, \dots, n\}). \end{cases}$$

Example ($G = P_2$)





We describe the fan Δ_{\square_G} of \square_G explicitly. Let

$$\mathcal{N}^\square(G) = \{N \subset \mathcal{I}_G \cup \{\{\bar{1}\}, \dots, \{\bar{n}\}\} \mid F_I \cap F_J \neq \emptyset \text{ for any } I, J \in N\}.$$

For $N \in \mathcal{N}^\square(G)$, we denote by σ_N the $|N|$ -dimensional cone $\sum_{I \in N} \mathbb{R}_{\geq 0} e_I$ in \mathbb{R}^n . Then $\Delta_{\square_G} = \{\sigma_N \mid N \in \mathcal{N}^\square(G)\}$.

Example

The fan $\Delta_{\square_{P_2}}$ of \square_{P_2} is illustrated in the right figure and thus the associated toric variety is $\mathbb{P}^1 \times \mathbb{P}^1$ blown-up at one point.



3. Toric Fano varieties

X : a nonsingular projective algebraic variety.

Definition

- X is **Fano** \Leftrightarrow the anticanonical divisor $-K_X$ is ample.
- X is **weak Fano** $\Leftrightarrow -K_X$ is nef and big.
 - A divisor D is **nef** $\Leftrightarrow (D.C) \geq 0$ for any curve $C \subset X$.
 - D is **big** \Leftrightarrow the litaka dimension $\kappa(X, D)$ is equal to $\dim X$.

Theorem

There are a finite number of isomorphism classes of toric (weak) Fano varieties in any given dimension.

dimension	1	2	3	4	5	6
# of toric Fano varieties	1	5	18	124	866	7622
# of toric weak Fano varieties	1	16	?	?	?	?

Let $\Delta(r)$ be the set of r -dimensional cones in Δ for $0 \leq r \leq n$.

Proposition

$$\Delta(n-1) \xleftrightarrow{1:1} \{\text{torus-invariant curves on } X(\Delta)\},$$
$$\tau \mapsto V(\tau).$$

Proposition

Let $X(\Delta)$ be an n -dimensional nonsingular projective toric variety.

- $X(\Delta)$ is Fano $\Leftrightarrow (-K_{X(\Delta)} \cdot V(\tau)) > 0 \quad \forall \tau \in \Delta(n-1)$.
- $X(\Delta)$ is weak Fano $\Leftrightarrow (-K_{X(\Delta)} \cdot V(\tau)) \geq 0 \quad \forall \tau \in \Delta(n-1)$.

Let Δ be a nonsingular complete fan in \mathbb{R}^n .

Proposition

Let $\tau = \mathbb{R}_{\geq 0}v_1 + \cdots + \mathbb{R}_{\geq 0}v_{n-1} \in \Delta(n-1)$, where v_1, \dots, v_{n-1} are primitive vectors in \mathbb{Z}^n , and let v and v' be the distinct primitive vectors in \mathbb{Z}^n s.t. $\tau + \mathbb{R}_{\geq 0}v, \tau + \mathbb{R}_{\geq 0}v' \in \Delta(n)$. Then:

- $\exists a_1, \dots, a_{n-1} \in \mathbb{Z}$ s.t. $v + v' + a_1v_1 + \cdots + a_{n-1}v_{n-1} = 0$.
- $(-K_{X(\Delta)} \cdot V(\tau)) = 2 + a_1 + \cdots + a_{n-1}$.

4. Main results

Let G be a finite simple graph.

Theorem 1 (S)

$X(\Delta_{\square_G})$ is Fano \Leftrightarrow each connected component of G has ≤ 2 nodes.

- $G = P_1 \Rightarrow X(\Delta_{\square_G}) = \mathbb{P}^1$.
- $G = P_2 \Rightarrow X(\Delta_{\square_G})$: $\mathbb{P}^1 \times \mathbb{P}^1$ blown-up at one point.

Hence if $X(\Delta_{\square_G})$ is Fano, then it is a product of copies of \mathbb{P}^1 and $\mathbb{P}^1 \times \mathbb{P}^1$ blown-up at one point.

Let G be a finite simple graph.

Theorem 2 (S)

The following are equivalent:

- $X(\Delta_{\square_G})$ is weak Fano.
- $\forall I \subset V(G)$, $G|_I$ is not isomorphic to any of the following:
 - (i) A cycle with ≥ 4 nodes.
 - (ii) A diamond graph (the graph obtained by removing an edge from a complete graph with four nodes).
 - (iii) A claw (a star with three edges).




Figure: a diamond graph.

Examples

- If G is a path or a complete graph, then $X(\Delta_{\square_G})$ is weak Fano.
- If G is a graph obtained by connecting more than two graphs with one node, then $X(\Delta_{\square_G})$ is not weak Fano.
- The toric variety associated to the graph cubeahedron of the graph below is weak Fano.




Figure: an example.

dimension	1	2	3	4	5	6
# of connected graphs	1	1	2	6	21	112
weak Fano	1	1	2	3	6	11

Thank you for your attention!

References

- M. Carr and S. L. Devadoss, Coxeter complexes and graph-associahedra, *Topology Appl.* 153 (2006), no. 12, 2155–2168.
- S. Choi, B. Park and S. Park, Pseudograph and its associated real toric manifold, *J. Math. Soc. Japan* 69 (2017), no. 2, 693–714.
- S. Choi and H. Park, A new graph invariant arises in toric topology, *J. Math. Soc. Japan* 67 (2015), no. 2, 699–720.
- C. De Concini and C. Procesi, Wonderful models of subspace arrangements, *Selecta Math. (N.S.)* 1 (1995), 459–494.
- S. L. Devadoss, T. Heath and W. Vipismakul, Deformations of bordered surfaces and convex polytopes, *Notices Amer. Math. Soc.* 58 (2011), no. 4, 530–541.

References

- T. Manneville and V. Pilaud, Compatibility fans for graphical nested complexes, *J. Combin. Theory Ser. A* 150 (2017), 36–107.
- T. Oda, *Convex Bodies and Algebraic Geometry. An Introduction to the Theory of Toric Varieties*, *Ergeb. Math. Grenzgeb.* (3) 15, Springer-Verlag, Berlin, 1988.
- B. Park, H. Park and S. Park, Graph cubeahedra and graph associahedra in toric topology, arXiv:1801.00296.
- H. Sato, Toward the classification of higher-dimensional toric Fano varieties, *Tohoku Math. J.* 52 (2000), no. 3, 383–413.
- Y. Suyama, Toric Fano varieties associated to graph cubeahedra, arXiv:1802.07059.